



SOLANA GmbH

Albert-Einstein-Ring 5
22761 Hamburg
Germany
Phone +49 (0)40 41 42 40-0
Fax +49 (0)40 41 77 16
info@solana.de
www.solana.de

Rossini



An early maincrop variety, suitable for crisps and dehydrated potato products. Its outstanding resistances against PCN Ro 1(9), Ro 3(9) as well as Pa 2(9) and Pa 3(9) make Rossini attractive for intensively farmed potato land. Next to PCN resistance, it is also resistant against potato wart disease 1. Tubers are round oval with a light yellow flesh colour. Good tolerances against bruising, blackleg and tuber blight. Irrigation is worthwhile. The highest crisping quality is achieved from greentop harvesting directly from the field until December.[nbsp] [nbsp]

Plant

Maturity	Medium early
Plant type	Stem type
Growth habit	Upright
Color of flower	Light red-violet
Foliage development	Rapid
Number of tubers	High
Dormancy	Medium, High

Tuber characteristics

Tuber shape	Round oval
Flesh color	Cream, Light yellow
Eye depth	Shallow, Medium
Skin color	Yellow
Skin characteristic	Netted

Resistances

Nematodes	Ro 1, Ro 3, Pa 2 (8), Pa 3
Potato wart disease	Race 1
Leaf blight	Medium, High
Tuber blight	Medium
Rhizoctonia	Medium
Black leg	High
Common scab	Medium
Internal rust spot	Medium
Damages	Medium
Bruising	Medium
PVY	Low
PVY ntn	High

Yield and Quality

Market outlet	Crisps, Dehydrated potato products
Yield	Medium
Share oversize	Low
Share undersize	Medium
After cooking discoloration	Low
Discoloration raw	Medium
Dry matter content	Medium
Chips quality after harvest	Good